

27 OCT 2004



INVESTOR IN PEOPLE



# **PRIORITY DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

REC'D 20 DEC 2004


WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Registration under the Companies Act does not constitute a new legal entity but merely affects the company to certain additional company law rules.



Signed

Dated 7 October 2004

BEST AVAILABLE COPY

# Patents Form 1/77

Patents Act 1977  
(Rule 16)



280CT03 EB47625-2 D02656  
P01/7700-0-00-0325019.8

## Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road  
Newport  
South Wales  
NP10 8QQ

1. Your reference

P15224 r2/sa

2. Patent application number

(The Patent Office will fill in this part)

0325019.8

3. Full name, address and postcode of the or of each applicant (underline all surnames)

GSF-Forschungszentrum für Umwelt und Gesundheit  
Ingolstaedter Landstrasse 1  
85764 Oberschleissheim  
Germany

Patents ADP number (if you know it)

8266690001

If the applicant is a corporate body, give the country/state of its incorporation

Germany

4. Title of the invention Floatable granular substrate for culturing plant material

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Hoffmann · Bitle  
Sardinia House  
Sardinia Street  
52 Lincoln's Inn Fields  
London WC2A 3LZ

Patents ADP number (if you know it)

07156466001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number  
(if you know it)

Date of filing  
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(day / month / year)


8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
  - b) there is an inventor who is not named as an applicant, or
  - c) any named applicant is a corporate body.
- See note (d))

Yes

## Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form	0
Description	42 ✓
Claim(s)	5 ✓
Abstract	1 ✓
Drawing(s)	7 + 7 

10. If you are also filing any of the following, state how many against each item.

### Priority documents

#### Translations of priority documents


Statement of inventorship and right to grant of a patent (*Patents Form 7/77*) 1 ✓

Request for preliminary examination and search (*Patents Form 9/77*) 1 ✓

Request for substantive examination (*Patents Form 10/77*) 1 ✓

Any other documents  
(please specify)

11. I/We request the grant of a patent on the basis of this application.

 Signature

Date  
27 October, 2003

12. Name and daytime telephone number of person to contact in the United Kingdom  
David Lethem  
020 7404 0116

### Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

### Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

# Floatable granular substrate for culturing plant material

## *Technical field of the invention*

- 5 The invention relates to a method for culturing plant material which makes use of a floatable granular substrate, and a culturing kit which comprises said substrate together with a culturing vessel.

## 10 *Background of the invention*

- The use of hydroponic systems for culturing plant material including whole plants, seeds, seedlings, meristems or calluses is widespread in science and industry. In particular, biotechnological methods of plant propagation, modification or culture typically involve hydroponic systems at least at some stage. Such biotechnological methods are typically used in science, but also are in routine use in generating plant material for industrial purposes, both in agriculture and the production of ornamental flowers or plants.

- Hydroponic systems are often chosen instead of conventional soil in order to provide chemically defined nutrients to plants. As a matter of fact, this is a technical necessity for many biotechnological propagation methods where defined cocktails of phytohormones need to be administered at certain points in the development of the plant. Furthermore, nutrient solutions are often easier to handle and to sterilize as compared to soil.



In scientific settings, hydroponic systems are often used in combination with phytotronic cabinets and exposure chambers supplying defined artificial light quality, quantity,  
 5 direction and temporal variation (Thiel et al, 1996). Together, these measures allow the creation of reproducible and defined growth conditions for the experiments.

Apart from growing plants under conditions of standardized  
 10 nutrient supply, hydroponic systems also allow simple addition of substances to the nutrient solution, without any such interferences that could be expected for plants cultured in soil. Further, plant material such as roots can readily be isolated from hydroponic systems for further analysis.

15 In hydroponic systems, plants are typically grown directly in granulated substrates such as expanded clay. Culture substrate is provided for mechanical support and an appropriate tactile environment for root growth.  
 20 Alternatively, in particular for scientific applications, highly specialized vessels, rockwool- or agar-based systems can be used (Gibeaut et al, 1996; Heidenreich, 1999; Huttner and Bar-Zvi, 2003; Tocquin et al., 2003).

25 However, such systems have considerable limitations:

---

In conventional hydroponic systems using granulated substrate, the level of the nutrient solution needs to be carefully adjusted to an optimal height in relation to the  
 30 substrate to provide optimal culturing conditions. If the nutrient solution is too high, the plant material may encounter detrimental anaerobic conditions. On the other hand, if the level of nutrient solution is too low, the plant

material may fall dry. This is of particular relevance for immature plant material, especially such without an established root system.

5 Also, granulated substrates often contain clefts that are too large for growing plants from small seeds, as these may easily fall into a cleft where they do not find optimal growth conditions. Therefore, seedlings are often germinated on one particular substrate (e.g. filter paper) and are then  
10 transferred into hydroponic culture. Such transferral can result in mechanical stress for the seedlings and furthermore is very labour and time intensive.

One improvement over hydroponic systems based on granulated  
15 substrates are floating culturing systems. Such floating systems can provide optimal growth conditions for differing levels of nutrient solution in the culturing container. One example of a floating system used in agriculture e.g. for the cultivation of tobacco seedlings contains a polystyrene-float  
20 with a number of openings, in which seedlings can be placed (Leal, 2001). However, each individual plant needs to be placed manually into one of the openings, hence this method is very labour intensive. Also, the seedling is supported in the opening by either soil, or a substrate like e.g.  
25 rockwool. These substrates are unsuitable for a number of applications.

Any system employing soil or other conventional substrates is limited by the fact that such substrate is not chemically  
30 inert. In particular, such substrates can readily adsorb components of the nutrient solution. This is a considerable disadvantage when the nutrient solution needs to contain a defined concentration of a specific ingredient. Such is the

case e.g. in many scientific experiments, or in biotechnological culturing methods wherein defined concentrations of phytohormones must be present at given growth periods.

5

Further to causing variation of concentrations of a desired ingredient, such adsorption by the substrate may necessitate the addition of larger quantities of the ingredient to obtain the desired final concentration. In case of expensive ingredients, such as many phytohormones, this can represent a substantial economic disadvantage, in particular for biotechnological methods of cultivation.

To overcome the limitations resulting from the adsorption of chemicals to the substrate, agar-based systems can be employed. However, these are expensive and laborious to set up. Furthermore, once the roots of the plant grow into the agar, the substrate cannot readily be exchanged. Hence, the temporally controlled addition or withdrawal of substances is limited. A further limitation of agar based systems is the adherence of the substrate to root material. When such a plant is replanted into e.g. soil for further growth, the agar residues are prone to bacterial degradation. This process of bacterial agar decomposition can result in unfavourable conditions for the plants. Further, residual agar attached to the plant material poses a technological limitation to e.g. experiments that study the uptake of certain chemicals by the plant and require the analysis of plant material absolutely free of surrounding substrate.

30

Alternative culture systems have been described. For *Arabidopsis thaliana* a floating sponge system for growth of individual plants has been used (Arteca & Arteca, 2000).

Aluminium tolerance of barley was tested with a 50 ml syringe system, which allowed the growth of 5 seedlings (Feng et al., 1997). However, both systems are laborious to handle, and cannot readily be scaled up. Hence, they cannot be used for industrial application or scientific experiments requiring large numbers of individual plants.

Osmotek Ltd., Rehovot, Israel, commercialises a floating culturing system ([www.osmotek.com/product.htm#LifeRaft](http://www.osmotek.com/product.htm#LifeRaft) and [www.osmotek.com/liferaftDescription.html](http://www.osmotek.com/liferaftDescription.html) ). In this system, a semipermeable membrane is placed inside a frame. This frame is then supported by a float that allows the membrane to be in contact with the nutrient solution. Float and frame are placed inside a proprietary culturing container filled with nutrient solution.

In particular for larger plant material this system is very laborious, as the plants must be placed in so called support sockets to prevent them from falling over, thereby losing good contact between the culture material base and the membrane.

The semipermeable membrane does not allow root penetration. Hence, there is no direct contact of plant and nutrient solution. This may affect both nutrient exchange and accumulation of toxic exudates. Alternatively, however, membranes with holes are available, that allow root growth into the nutrient solution.

However, if plant material is grown through holes in the membrane, it is very difficult to isolate seedlings from the membrane without destruction of root tissue. Also, such membranes cannot readily be reused.

Even if the membranes have holes, contact of plant material with the nutrient solution is restricted by the float. Though the float contains a central opening to allow solution  
 5 exchange, all regions not directly apposing this opening only enjoy a very thin solution support of a few millimetres. The practical consequence of limited or missing contact with a large solution phase is a limited nutrient supply on the one hand, and accumulation of toxic plant exudates on the other  
 10 hand.

This culturing system brings about a number of further limitations. For example, the user is restricted to specialized culturing containers, that firstly are expensive,  
 15 and secondly cannot readily be scaled up, as the integrated system is only available in few defined sizes. Furthermore, any float is laid out to support a defined plant weight. If the plant weight differs from that weight, it can become necessary to change the float.

20

A further limitation of the membrane system is that primary roots may not receive the optimal tactile signals necessary for development. This is of particular importance, if the plant material is meant to be replanted for further  
 25 cultivation, or if entire plants of normal morphology are needed for scientific purposes.

---

Accordingly, there is a need for a culturing system that overcomes all of these limitations. In particular, the  
 30 culturing system should readily be scaleable from laboratory- to industrial scale, should be easy to set up without laborious preparation, should be made of cheap components and allow use of various culturing containers. Additionally the

system should allow culturing of different plant materials, including seeds of different sizes, seedlings, plants, meristems, calluses or other cellular aggregates. In particular, the culturing system should allow culturing of  
 5 plant material bearing roots, and should allow easy harvesting and replanting of such material, it should provide plant material with sufficient nutrients and dissipate toxic exudates over a prolonged period of time, should allow easy exchange of culture medium, it should be chemically inert not  
 10 to adsorb ingredients from the culture medium and should be sterilizable by conventional means.

# *Summary of the invention*

The present invention provides the use of a floatable  
 15 granular substrate for culturing plant material.

Preferred embodiments of said floatable granular substrate comprise particles having an average diameter in the range of 1-25 mm, more preferred 1-10 mm, a regular or irregular  
 20 spheroidal or polygonal shape and a smooth surface.

Another preferred embodiment comprises a granular substrate that is chemically inert.

25 Another preferred embodiment comprises a granular substrate with a density of 50 - 99.8 % of the density of the culture medium used, or a density in the range of 0.5 - 1.1 g/cm<sup>3</sup>.

Another preferred embodiment comprises a granular substrate  
 30 being a thermoplastic polymer comprising one or more of high density polyethylene (HD-PE), low density polyethylene (LD-PE) or polypropylene (PP).

Another preferred embodiment comprises particles composed of more than one component, wherein said components can individually be more or less dense than the average density  
5 of the particle, and / or particles comprising at least one hollow enclosure.

Another preferred embodiment comprises a granular substrate sterilizable by a chemical treatment, irradiation and / or  
10 heat.

In one preferred embodiment, the granular substrate forms a floatable substrate layer with a thickness of 0.5 - 20 cm, and preferably 0.5 - 10 cm that may or may not float on a  
15 culture medium which may or may not be aerated.

In another embodiment the granular substrate layer comprises embedded additional support structures.

20 According to a further aspect, the present invention provides a culturing kit for culturing plant material comprising a floatable granular culture substrate. Said substrate can be combined with any one, two or all of the following additional components: culturing solution, plant material and culturing  
25 vessel. Granular substrates of all previously specified embodiments can be used for said kit.

According to a further aspect, the present invention provides a method for culturing plant material comprising:

- 30 (a) forming a layer of floatable granular substrate in a culturing vessel,

- (b) placing plant material on or in said layer, and
- (c) culturing the plant material in the presence of a culture medium, wherein there is no additional structure supporting the plant material from underneath.

5

The method can be performed using any of the previously specified culturing kits or any of the previously specified embodiments of the granular substrate.

10

#### ***Brief Description of the Figures***

**Figure 1:** Cultivation of barley seedlings on a floating layer of LD-PE substrate.

15

**Figure 2:** Representative example of (a) six day old barley seedling harvested from LD-PE substrate, and (b) tobacco derived meristems showing rich primary root development harvested from PP substrate after 10 days of culturing.

20

**Figure 3:** Example of calluses cultivated on floating LD-PE substrate (a) a lateral view, and (b) a top view.

**Figure 4:** Example of meristems derived from tobacco on floating PP substrate in a glass culturing vessel. (a) a lateral view and (b) a top view.

25



*Detailed Description of the Invention*

**Substrate Material**

Conventional substrates such as expanded clay, rockwool, soil or agar are not floatable. If floating is desired, such  
5 substrates must be suspended by a separate floating entity. In contrast, the present invention provides a floatable granular substrate for culturing plant material.

The material for the substrate according to the present  
10 invention is not limited, as long as this material is floatable in a culturing solution, and provides mechanical stability and tactile stimuli for root growth in a hydroponic system. The elasticity of the substrate material is not  
limited. Hence, any materials ranging from highly stiff to  
15 highly elastic can be used. Preferably, the material is chemically inert. Examples of substrates with these characteristics comprise, but are not limited to, thermoplastic polymers of appropriate density. Specific examples of such thermoplasts comprise polypropylene (PP),  
20 high density polyethylene (HD-PE) or low density polyethylene (LD-PE). These polymers are used in large quantities in a wide range of industrial applications and are typically provided in the form of a raw material granulate. Therefore, such granulate can readily be obtained at low cost and in any  
25 reasonably desired quantity. The granulate can be directly  
used as a culture substrate, or be further processed by e.g. selecting a certain size fraction, sterilization, packaging, etc.

30 For particular applications, a preferable granulate material may be degradable in a defined way. Examples of such materials comprise, but are not limited to e.g. extruded

starch polymers. It may be required to coat such degradable materials in an appropriate fashion to achieve the desired properties. It is obvious to the skilled person, what kind of degradable composition to use for a given application.

5

In the present specification, the term "culturing" of plant material is used in its widest sense. This means that plant material is generally kept under such conditions which the skilled person expects the plant material will survive at.

10

However, that does not exclude that under certain circumstances conditions will deliberately be chosen such that the plant material does not survive. Culturing can, but need not, mean that plant cells divide or differentiate.

15

Examples of such processes include growing seedlings from seeds, differentiating calluses, or cultivating meristems.

20

In the context of the present invention, the terms "plant" and "plant material" are used in their widest senses and are meant to include all conceivable elements derivable from a plant, such as organs (e.g. roots, leaves, etc), tissue isolated thereof (e.g. meristems), or individual cells or cellular aggregates (e.g. calluses) isolated thereof. It further comprises all stages of plant development, comprising but not limited to seeds, seedlings or fully grown plants.

25

The term "granular" means that the substrate is composed of a multitude of individual particles. Said particles are not firmly attached to one another. Individual particles can be of differing size or shape within a certain degree of freedom, such that a given culture substrate comprises a multitude of similar particles.

30

For the present invention, the precise shape of the granules is not limited, but will preferably approximate to a spheroidal shape or a polygonal shape. Other examples of possible shapes comprise, but are not limited to regular or irregular polygonal shapes, cubes, octahedra, tetrahedra, pyramids etc. In this context, "spheroidal" means that the particles of the granulate have a globe or ball shape in the broadest sense. This does not preclude that individual particles or the majority of the particles of the granulate approximate to the shape of an ideal sphere.

Typically, however, the particles of the predominant shape will deviate from the ideal geometrical shape and bear some kind of irregularity. This means e.g. that particles can be ball shaped, ellipsoid, pear shaped or any other shape that in the widest sense can be considered spheroidal. Furthermore, the particles can contain clefts, ridges, corners or edges, one or a few spikes or one or a few holes.

In other words, the overall appearance of the particles preferably resembles spheres or polyhedrons to some degree. This is a requirement to form floatable layers of sufficient stability to carry the weight of plant material. The irregularity of shapes and the size distribution in the granulate will have an impact on packing density of the substrate. The more individual particles approximate to ideal spheres and the more homogenous the size distribution, the more the packing will approximate to the theoretical optimum of tightly packed spheres.

Though the shape of individual particles can be irregular, the surface of the particles should be smooth. This does not preclude the presence of one or a few clefts, ridges, corners

or edges, one or a few spikes or one or a few holes. In particular, the substrate should not have any extensive surface protuberances or cavities that could comprise a solution phase separated to a substantial degree from the surrounding solution. Hence the substrate should be non-porous, i.e. it should not contain extensive inner surfaces and cavities that communicate with the outside. This does not preclude the presence of one or a few such pores and cavities, in particular if the diameter of their opening to the surrounding solution is relatively large in relation to their volume. If a larger number of pores with relatively small openings is present, they might form reservoirs for the diffusion-exchange of substances with the main solution body. Such inner surfaces are not preferred in the context of the present invention, but are typical for expanded clay, for example, and may prohibit the controlled temporal addition or withdrawal of substances, in particular when such substances are present in trace amounts.

The colour and transparency of the particles is not limited in the present invention. It can range from transparent to opaque and from black to white, including any chosen colour. For specific applications, transparent particles will be preferred, or such particles that have defined optical properties. Transparent particles may allow the optical investigation of the rhizosphere. If light of certain wavelengths, e.g. laser light, fluorescence of certain wavelengths should be used in such investigations, the granulate must have suitable optical properties that can readily be chosen by the skilled person.

The exact size of individual granules is not limited in the present invention, as long as it facilitates formation of a

floatable substrate layer of adequate stability. A preferred size range comprises average diameters in the range of 1 - 25 mm. In a typical application, the size- and shape- distribution of commercially available industry grade polymer granulate, such as PE or PP granulate, is suitable to form a substrate layer according to the invention. Granulate that contains irregular, round shapes with an average diameter ranging from 1 - 8 millimetres is preferably used. For specific uses, particle sizes representing a fraction of this range, e.g. 1 - 3 mm or 2 - 5 mm may be particularly preferred.

The granular structure of the substrate used in the present invention will result in the formation of clefts in the substrate layer, which will be filled with nutrient solution. The degree of cleft filling depends on capillary effects that are influenced by the size and shape of the granules. Further, cleft filling is influenced by the relative density of the substrate in relation to the nutrient solution.

20

The density of the granular substrate must be such that it allows the substrate to float on standard nutrient solutions (nutrient solution, culturing solution and medium are used synonymously). Thus, the density of the substrate will be lower than that of the solution. At the same time, it must not be so low that the substrate layer sits on top of the solution in such a way that the clefts between the substrate granules are not filled with solution. Therefore, an optimum relative substrate density in relation to solution density exists. This optimum is within the range of 99.8-50% of solution density.

30

In case the granulate comprises a composition of more than one substance, the mean density of the final granulate as used needs to fall within this range. Hence, single components of such a composite granule may have densities that do not fall within the specified ranges. For example, it can be envisaged that individual particles of the substrate may comprise gas filled enclosures, such as materials with a closed foam structure. In this case, the particle may comprise wall material with a density that may significantly exceed the density of the culture medium. Alternatively, particles may be comprised in part by high density material and in part by low density material. Materials that may be used for composite particles comprise, but are not limited to, polyvinylchloride (PVC), polymethylmetacrylate (PMMA), polytetrafluorethylene (PTFE), polycarbonate (PC), polyisoprene (PI), polyamide (PA), polyisobutylene (PIB), polyurethane (PU) and polystyrene (PS).

For a typical application, it is important that there is no absence of solution filling in a significant upper segment (e.g. a quarter of the substrate layer) of the layer. Also, the substrate layer should for a typical application not be immersed such as to be covered by a closed layer of solution, as it is desired to grow plant material positioned at the interface of solution and air.

---

However, for specific applications, such as studying the effect of draught or flooding, the density of the culture substrate may be chosen such that either a significant layer of substrate devoid of culture solution exists, or the substrate is fully immersed.

Many culturing solutions have a density approximating 1 g/cm<sup>3</sup>. Specific examples of such solutions are e.g. Hoagland's E-Medium, ASTM STP 1027, or MS-basal medium (available from Sigma; M5519). For such nutrient solutions, 5 typically the density of the floatable granular substrate will be in the range of 0.5 - 1.00 g/cm<sup>3</sup>. A preferred density range is between 0.90 - 0.96 g/cm<sup>3</sup>. Specific examples of granular substrate in the desired density range are high density polyethylene (PE-HD, 0.94 - 0.96 g/cm<sup>3</sup>), low density 10 polyethylene (PE-LD, 0.914 - 0.928 g/cm<sup>3</sup>), or polypropylene (PP, 0.90 g/cm<sup>3</sup>).

However, for special applications where nutrient solutions of higher density are used, such as e.g. studying plant growth 15 under high salt concentrations, the density of the culturing substrate may be in excess of 1 g/cm<sup>3</sup>, e.g. 1.1 g/cm<sup>3</sup>.

Hence, the absolute density ranges of the substrate specified are not limiting. Rather, under any specific condition 20 chosen, the optimum proportion of densities of solution and substrate must be conserved to ensure the culturing substrate is floatable on the specific solution used.

It is important in the context of the present invention that 25 the floatable culturing substrate can provide such support from the beginning of the culturing process to the chosen plant material, and in particular small plant material such as certain seeds, that the plant material is prevented from sinking too deeply into the culture solution. Hence, any 30 substrate that requires additional mechanical devices at least at some point during the culturing process that support the plant material from below to prevent its sinking is not a

culturing substrate in the sense of the present invention. For example, this applies when plant material is supported by a horizontal structure that allows plant access to nutrients like a mesh, membrane or filter, and then is fixed onto this horizontal structure by adding some kind of substrate from above. Such a covering layer is not a culturing substrate in the sense of the present invention.

An additional feature that is desirable for some applications is that the substrate can readily be sterilized by conventional means, such as irradiation (gamma-irradiation, UV, etc.), chemical treatment, autoclaving or any combination thereof. For example, the substrate can be sterilized by a chemical treatment using any conventional chemical sterilizing agent. Such agents can contain e.g. halogens or halogenated compounds (Cl, I, Br, F), lower alcohols (e.g. ethanol, propanol), phenols or phenol derivatives, aldehydes (e.g. formaldehyde, glutaraldehyde), quaternary nitrogen compounds, amphoteric compounds liberating reactive oxygen species (e.g.  $H_2O_2$ ), NaOCl, or ethyleneoxide. Alternatively, the substrate can be gamma-irradiated. Alternatively, the substrate can be autoclaved at conditions known to the skilled person. A typical autoclaving temperature would be up to  $130^{\circ}C$ . In the latter case, only such substrate materials can be used that have a melting point higher than  $130^{\circ}C$ . For specific applications readily known to the skilled person, higher or lower autoclaving temperatures may be used. Thus, two groups of substrate materials can be distinguished based on their temperature stability, a low- and high-temperature stable group. The low temperature group has a thermal stability of maximally  $130^{\circ}C$ , the high-temperature group in excess of  $130^{\circ}C$ . Specific examples of low-temperature stable substrates are LD-PE (melting point:  $60-75^{\circ}C$ ) and HD-PE



(melting point: 90-120°C). A specific example of a high-temperature stable substrate is PP (melting point: 140°C).

**Chemical Properties:**

5 Substrate materials that are meant to be chemically sterilized must have a chemical stability sufficient to resist such treatment. Specific examples of such materials are LD-PE, HD-PE or PP, all of which have a chemical stability that allows chemical sterilization with typical  
10 agents used for this and known to the skilled person.

Apart from chemical stability that allows sterilization, it is preferable for the present invention that the chemical properties of the substrate are defined and controllable. In  
15 particular, it is preferable for the culture substrate of the present invention to be chemically inert. This means that the substrate should not readily interact with typical ingredients of plant nutrient solutions. In this context, to interact means the compound is adsorbed, precipitated,  
20 catalysed, oxidized, reduced, cleaved or chemical groups are added when in contact with the culture substrate. Adsorption in this context means that a physiologically relevant proportion of an added substance adheres to the substrate. This is of particular relevance when the substance is present  
25 at low concentration. In a most preferred embodiment of the present invention, the substrate does not readily adsorb ingredients of the medium. Typical ingredients of culture media comprise macronutrients such as phosphorus or nitrogen, micronutrients such as various metals, as well as  
30 phytohormones. Further, such substances that are typically used in physiological experiments in a scientific setting should not interact with the substrate. These include any xenobiotics such as heavy metals, inorganic compounds,

organic compounds, etc., but also comprise amino acids, peptides, proteins, nucleotides, RNA, DNA etc. The list of substances listed here is not exhaustive, but merely represents a selection of examples to illustrate the scope of the chemical space to which the substrate should essentially be chemically inert.

Particular requirements related to chemical inertness are that the culture substrate should not have any free reactive groups, and should not carry a surface charge or exposed charged residues. These properties should be stable over a wide pH range as might be used in culturing plant material. This pH range is from acidic (pH=1) to alkaline (pH=13). Specific examples of materials fulfilling these criteria sufficiently to be useful in the context of the present invention comprise, but are not limited to, PP or PE.

Defined and controllable chemical properties in the context of the present invention may mean for specific applications that chemical reactions, including interactions with the medium or release of compounds are desired.

For example, the granulate can comprise a functional chemical compound, such as a reactive covering. Such covering may contain a pH indicator or a temperature indicator which gives rise to visual information concerning the culturing conditions. Alternatively, the functional chemical compound comprised in the granulate may be any chemical of a defined, desired function, such as chemicals bearing e.g. fungicidal, algacidal, or microbicidal function. Such chemicals may also comprise substances that function as nutrients or toxins for plant material, or have any other measurable effect on plant

material. In some applications, the controlled and sustained release of such compounds will be desirable.

### Culturing vessel

5 To culture plant material according to the present invention, the culturing substrate will be added to a culturing vessel (the terms vessel and container are used synonymously). The container used for the invention is not limited in size or shape, as the granular substrate will readily form a closed  
10 substrate layer independent of container shape or size. Any container from laboratory scale such as glass or plastic tubes, beakers, bowls or troughs up to industrial scale growing chambers or troughs in any desired size, even in the hectare range, can be used. In the case of very large  
15 containers, application of the substrate will require appropriate mechanical means.

The combination of cheap and readily available culture substrate and the possibility of using a wide range of  
20 culture containers results in a culturing system easily scalable from scientific up to industrial scale. This scalability is independent of expensive investment in e.g. specialised culturing containers and hence provides a significant benefit to the present invention.

25

### Aeration of culturing solution

If required, gases such as air or any other gas can be applied to the culturing solution. For this purpose, an opening in the floating substrate layer is introduced. Such  
30 an opening can, for example, be introduced by a cork ring with its opening being free of culture substrate. Many other

ways of physically creating a substrate free area can readily be envisaged. The desired gas can be applied to the solution by suitable physical means in the area free of substrate. Such means comprise any kind of gas outlet that can be introduced into the solution such that the gas is set free in the region free of substrate.

### Culturing Kit

Though the specific vessel used for the present invention is not limited, according to an embodiment of the invention, the floatable culture substrate and an appropriate culture vessel form parts of a culture kit. Such a culturing kit may further include an appropriate culture medium, and possibly also plant material. Different culturing kits can be specifically adapted to a particular application. As an example, a culture kit may comprise a decorative vessel, containing culturing substrate comprising seeds of ornamental plants, and a separately packaged culture medium that simply needs to be added in order to start the culturing process. Alternatively, the culture kit may just comprise the floatable substrate and separately packed substrate-seed mixture, etc. In this way, the present invention provides a diversity of culture kits for hobby, science, or industrial use.

### —25— Culturing process—

According to the specific requirements of the plant material to be cultured, any of the following culturing steps may be combined with appropriate measures of sterilization and / or aseptic handling. Sterilization measures for media, containers, substrate, nutrient solution and plant material are widely known in the art. It is obvious to the skilled

person which kinds of culturing steps require such measures, therefore they will not be explicitly mentioned in the following description of the culturing method according to the present invention.

5

Typically, the floatable granular substrate will be added to the culturing container to form a layer of culture substrate. Plant material may either be added to the granular substrate prior to layer formation, or thereafter. A sufficient

10

quantity of substrate chosen to support the size and weight of plant material of interest can readily be determined. In this context it is important to note that the substrate layer is floatable, yet, it need not float at all times during the culturing process. It can be envisaged that at selected time-

15

periods, the substrate layer is either floated or rests on the bottom of the culture container. As an example, seedlings may be grown from seeds on a floating layer to allow for

optimal nutrient solution supply. Given that the weight of the plant material is relatively small in comparison to the

20

weight of the substrate layer, it will not significantly affect the degree of immersion of the substrate layer. Hence, cleft filling will be optimal for a wide range of plant material weight.

25

Still, at a later stage, when the seedlings have grown bigger and become too heavy to be supported by the floating layer, the layer may be lowered to the bottom of the container.

Then, the seedlings and in particular their roots have developed sufficiently that small variations in solution

30

height within the substrate layer are not as detrimental as at the beginning of the culturing process.

Hence, the culturing method employing a floating layer of culturing substrate is restricted to plant material of appropriate size and weight which permits suspended growth. Obviously, the culturing system may e.g. not be suitable for  
5 very large and heavy seeds such as certain nuts, but may easily support fully grown plants of small species.

However, to adapt the method to growing larger plants, additional support structures embedded within and supported  
10 by the floatable granular layer may be present. Such structures may e.g. be introduced to increase the mechanical stability of the overall layer, e.g. when culturing particularly large and heavy plant material. Such structures may define enclosed spaces within the granulate layer,  
15 wherein individual granules are still freely movable. In this context "enclosed" does not mean that physical barriers that substantially restrict medium exchange exist. If a multitude of such structures is present, the spaces between the enclosures is filled by un-enclosed granules. Examples of  
20 suitable structures comprise plastic nets with a mesh size smaller than the average granule size, or grids that are interposed in the layer. Bags formed from net material may thus enclose a portion of granules that are part of the layer. In contrast, grids may have considerably larger  
25 openings than the average granule size and yet increase the mechanical stability of the layer they are embedded in. In case the embedded structures define enclosures containing portions of particles, such enclosures can be used to separate the plant material attached to them from the rest of  
30 the layer, e.g. for harvesting and / or replanting.

Alternatively, the substrate layer may contain agglomerates or aggregates of substrate particles of different sizes,

wherein individual particles are attached to other particles to a variable degree and with variable attaching strength. Such structures would be interspersed with unattached particles to form a dense substrate layer.

5

The thickness of the layer of the floatable granular substrate is not limited, as long as the layer provides sufficient stability and buoyancy for the chosen application. Obviously, the thickness should be adjusted to the size and weight of the plant material to be cultured. The thickness can be adjusted either at the beginning of the culturing process by adding the appropriate amount of substrate material to the culturing vessel, or can be adjusted during the culturing process e.g. by adding substrate particles from below in a suitable culturing vessel. Hence, the present invention allows the continuous adaptation of the substrate layer thickness to optimal culturing conditions without any disruption of the plant material.

20 In a typical application growing plant material such as seedlings, meristems or calluses, the thickness of the substrate layer will be in the range of 0.5-20 centimetres, preferably 1-10 centimetres, more preferably 2-8 centimetres. Apart from the thickness of the layer, the average size of the culture substrate granules can be adjusted to the size and weight of the plant material to be cultured.

When using small seeds, such as e.g. derived from *Arabidopsis* or *Thlaspi*, an average granule size of 1-2 millimetres and a layer thickness of 2-3 centimetres are recommended. For larger seeds, such as e.g. barley, an average granule size of up to five millimetres and a layer thickness of approximately 5 centimetres is suitable. For even larger or heavier plant

material, such as large seeds, calluses or meristems derived from tobacco, lettuce, or tomatoes, layer thickness may be increased up to approximately 8 centimetres. Obviously, culturing of very large plant material will require layer heights in excess of 8 centimetres. There is no theoretical upper limit of layer thickness, other than the restrictions imposed by the culturing vessel.

When the granular substrate has been distributed in the culturing container to the desired layer thickness, plant material may simply be placed on top of the substrate layer or can be stuck into the layer to a desired depth.

An alternative to placing plant material on top of the substrate layer is mixing it with substrate. This is a preferred method when e.g. culturing seedlings from seeds. The seeds can be mixed with an excess of culture substrate and the resulting mixture can be distributed in a thin layer on top of the substrate layer. The seeding density can readily be adjusted by choosing different ratios of seeds / culture substrate. A typical ratio would involve e.g. approximately a ten-fold volume of culture substrate compared to seeds.

The simple preparation of the substrate layer by distribution in an appropriate culture container, in connection with the easy application of plant material represents a significant advantage over many conventional systems. Hence, laborious substrate preparation such as e.g. in an agar-based system, floating sponges etc. is not required. Also, the often most laborious step, the correct positioning of plant material, is greatly facilitated. This is particularly obvious in comparison to conventional floating systems, where plants need to be placed in or on top of individual openings. It can



be envisaged, that the present system will allow machine-planting in an industrial setting, where at present plants are positioned by hand.

- 5 Any appropriate culture medium can be used for the present invention, given that the density of the granular substrate is chosen accordingly. A wide variety of different nutrient solutions are in routine use in hydroponic plant culture. Examples of such solutions comprise, but are not limited to,
- 10 Hoagland's solution or MS-basal medium.

The nutrient solution is either added by pouring it in at an edge of the container, or it can be flooded from the bottom of the container, depending on the technical specification of

15 the culturing container. Obviously, culturing vessels can also be constructed such that a permanent or intermittent flow of medium through the vessel and the substrate layer therein occurs. Such flow can be used to achieve defined

20 temporally defined addition or withdrawal of certain substances, or the complete exchange of medium. Such flow may also contribute to providing sufficient medium of a precisely defined composition and physical parameters to the plants.

- 25 In a particular application, flow through systems can be employed e.g. to establish an online indicator system in e.g. a waste water stream, wherein a readily detectable change in the culturing system would indicate a defined environmental condition. Such flow through may also be employed when using
- 30 the present invention in test systems other than online sensors.

When medium is added, initially, the clefts of the culturing layer will fill with solution. Then, if more solution is added, the substrate layer will begin to float. The height of the solution phase under the substrate layer depends on the amount of solution added. The height of the solution phase can freely be chosen according to the specifications of the container and the desired culturing conditions.

For some applications, such as e.g. further culturing of seedlings or plants that already have a developed root system, it may be advisable to plant on a floating substrate layer, rather than before nutrient solution is added. Of course, all other plant material can also be added at this point rather than before solution is added.

Now, plant material will grow in the hydroponic system. Roots will find both sufficient physical stability to support growth, and the necessary tactile stimuli for development of primary roots. In case the plant material develops roots and is grown for a sufficient length of time, and given that the culture substrate is floating, the roots may grow out of the substrate layer into the nutrient solution.

Hence, an advantage over e.g. available membrane based culturing systems is that root development is not restricted by any solution / membrane interfaces.

If the culture substrate is floating, the excess of nutrient solution will provide sufficient water and nutrients for the plant material over a prolonged period of time, and will dissipate toxic exudates. This represents an advantage over all systems that contain areas of restricted nutrient solution volume, such as commercially available membrane

based floating systems. Further, the solution-air interface will remain unchanged at the upper edge of the substrate layer for a wide range of solution volumes. This is an advantage over conventional, non-floating systems..

5

If necessary, the culturing solution can readily be exchanged during the culturing process, without physically disrupting the plant material. Thus, for example, different nutrient solutions can be used for different growth phases of the plant material. Such an exchange is impossible in soil-grown plants, but also in e.g. agar-based systems. Also, hydroponic systems employing e.g. expanded clay may exhibit a substantial retention effect due to the extensive inner surfaces of that substrate.

15

To harvest plant material, it can simply be pulled out of the substrate layer, without causing any physical disruption to the plant material. This is a requirement both in industrial applications, where the plants are cultivated further, and in science, when intact plant material needs to be harvested. In comparison, if plants grow through membranes, they need to be carefully excised. Also, any systems employing rock wool or agar make it very difficult to separate roots from substrate.

20

25

Further, such systems do not allow to readily replant in a different culturing container, in particular where a root system has developed. Non-floating hydroponic systems require emptying of the substrate. Agar-based systems require excision of individual plants. The remaining agar may produce rot if transferred onto e.g. soil as a substrate for further growth. Systems where plants grow through holes in a supporting structure can result in mechanical damage to plant

30

material, or the material may even get stuck when growing for too long.

Consequently, all these systems are labour intensive in replanting, and some do not allow replanting of root bearing plants at an industrial scale at all. In contrast, when using a floating, granular substrate layer, plants can easily be harvested and easily be replanted. For replanting, the plants can simply be gently pushed through the layer of floatable substrate, until the roots can spread freely in the solution phase, and the body of the plant is firmly anchored in the substrate.

Thus, the floatable granular substrate of the present invention may facilitate automated harvesting and / or replanting devices that cannot readily be used on conventional systems.

Further, it is possible to add a layer of conventional substrate like soil at a given point in the culturing process, to allow the plant material to grow into said conventional substrate. Such a process may e.g. facilitate and speed up consecutive replanting and hardening.

Previously, a culturing process employing PE beads has been described (Hart JJ et al., 1998 a and b). In this process, seeds were germinated on filter paper. Then, seedlings were placed on mesh bottom cups and covered with black PE beads to shield them from light. The cups were suspended in a vessel containing medium. Thus, the PE beads in this process are not a culturing substrate in the sense of the present invention, as they do not provide sufficient support from underneath to prevent the plant material from sinking. Rather, the plant

material is supported by the mesh bottom of the culturing cups. The beads are functioning as a mere cover that weighs the seedlings down on the mesh bottom and shields them from light. Further, the PE beads are not a culturing substrate in the sense of the present invention as they do not allow germination of seeds and culturing seedlings without repositioning. Also, the seeds are not meant to float, quite the contrary, the beads were used as weights to firmly position the seedlings on the mesh bottom. Finally, in this culturing process the mesh bottom forms a mechanical structure that permanently separates plastic beads and free solution phase throughout the culturing period. The culturing process of the present invention does not require such a mechanical barrier.

### Examples

#### Culture medium

MS-basal medium (Sigma) was preferably used for culturing plant material in the consecutive examples. Alternatively, Hoagland's E-Medium was prepared as follows (additions per liter of final medium):

Substance	Stock solution (g/100ml)	Addition to final medium (ml/l)
MgSO <sub>4</sub> •7 H <sub>2</sub> O	24.6	1.0
Ca(NO <sub>3</sub> ) <sub>2</sub> •4 H <sub>2</sub> O	23.6	2.3
KH <sub>2</sub> PO <sub>4</sub>	13.6	0.5
KNO <sub>3</sub>	10.1	2.5
Micronutrients (see below)		0.5
Fe-EDTA Solution added last (see below)		20.0

Adjust pH to 5.8 with NaOH or HCl. Sucrose may be added as 10 g / l if the culture is axenic. Autoclave.

**Micronutrient Stock Solution:**

5		
	H <sub>3</sub> BO <sub>3</sub>	2.86 g/l
	MnCl <sub>2</sub> ·4 H <sub>2</sub> O	1.82 g/l
	ZnSO <sub>4</sub> ·7 H <sub>2</sub> O	0.22 g/l
	Na <sub>2</sub> MoO <sub>4</sub> ·4 H <sub>2</sub> O	0.09 g/l
10	CuSO <sub>4</sub> ·5 H <sub>2</sub> O	0.09 g/l

**Fe·EDTA Stock Solution:**

	FeCl <sub>3</sub> ·6 H <sub>2</sub> O	0.121 g/250ml
15	EDTA	0.375 g/250ml

Dissolve completely and make up to 250 ml. After autoclaving medium, add Fe·EDTA Stock Solution aseptically.

20

**Culturing conditions**

In the examples the following growing conditions were used unless specified otherwise: Plant material was cultivated for 6 d to 14 d in a controlled environment cabinet (relative humidity 70 +/- 5 %) under a 14/10 h day-night cycle with a photosynthetic active radiation of  $116 \mu\text{M}\cdot\text{m}^{-2} \text{ s}^{-1}$  from 06:00 to 20:00 h middle European summer time and a temperature of 24/20 °C.

30

**Example 1: Culturing barley on floating culture substrate**

**Plant material:**

*Hordeum vulgare* cv. Barke was used. Cv: Barke (BSA-Nr. 1582) is an awned, double-lined summer barley, which were breded 1996 by Breun from Libelle X Alexis.

5 **Pre-treatment of barley caryopses under axenic conditions:**

Caryopses were incubated for 1 min in ethanol (70 vol / vol %) and then for 1 min in H<sub>2</sub>O distilled. Caryopses were then incubated two-times for 5 min in NaOCl (10 vol / vol % of a stock solution containing 6-14 % free, active chloride),  
10 containing 0.1 vol / vol % Triton X-100. The caryopses were then washed ten-times for 1 min with H<sub>2</sub>O distilled and swelled for 24 h in H<sub>2</sub>O distilled at room temperature.

**Floatable substrate layer**

15 LD-PE-granulate with an average granule size of 3-5 millimeters was used. A layer of granulate of approximately 4 cm in height was poured into different glass beakers. The beakers ranged in size from 50-5000ml, with a corresponding diameter of approximately 4 - 25 centimetres. Alternatively,  
20 rectangular polyethylene troughs with a size of approximately 50x70 centimetres and a height of approximately 10 centimetres were used. The required volume of granular substrate was 40 l / m<sup>2</sup>. The caryopses from Example 1 were mixed with PE-granulate (1:10; v/v). This mixture was  
25 distributed on top of the PE-granulate layer, resulting in a seedling density of 10-15 plants per 10 cm<sup>2</sup>. Alternatively caryopses were applied directly onto the PE-granulate.

Then, Hoagland's medium or MS-basal medium was added by  
30 pouring into the culture container at the edge. Pouring speed was chosen such as not to upset the substrate layer. The solution volume was chosen such that a liquid phase of

approximately 4 cm was formed underneath the floating substrate layer.

Seedlings were grown as indicated. Thereafter, seedlings were harvested by gently pulling them out of the substrate layer. Representative seedlings obtained by this method are depicted in figure 1 and 2a. The seedlings had reached an average height of 100-150 mm and an average weight of up to 2g, and showed a normal morphology. These values correspond well with such known from the literature for respective seedlings grown on e.g. agar substrate.

Hence, floatable substrate is capable of supporting normal growth of barley seedlings in large quantities under strictly controlled culture conditions.

#### Example 2: Comparison of heavy metal uptake of barley seedlings on floating culture substrate or agar substrate.

20

MS-basal Medium or Hoagland's medium was supplemented with heavy metals as follows: stock solutions of  $\text{HgCl}_2$  (0.1 M) or  $\text{Cd}(\text{NO}_3)_2 \cdot 4 \text{H}_2\text{O}$  (0.1 M) were applied to the autoclaved medium when it had cooled to about 55 °C such that final heavy metal-ion concentrations of 10, 20, 30 and 40  $\mu\text{M}$  were obtained.

25

Agar culturing substrate was prepared by adding 1.5 wt / vol % agar (Sigma) to Hoagland's or MS-basal medium. After heating to 100 °C, respective amounts of heavy metal salts were added when the medium had cooled down to 55°C. Then the

30



mixture was poured into Phytotray II™ culture vessels (Sigma) and left to cool for subsequent use.

Barley caryopses prepared and seeded onto a floatable substrate layer as in Example 1. Alternatively, the caryopses were directly placed on culturing agar as described above.

Seedlings were grown as described, and leaves and roots of 6 day old plants were harvested and weighed to determine the fresh weight. Samples were shock-frozen in liquid nitrogen and stored at -80 °C.

Total mercury and cadmium content of leaves and roots was determined by electrothermal vaporization inductively coupled plasma-mass spectrometry (Michalke et al., 1997).

The results are shown in Table 1:

Table 1: Concentration of heavy metal ions ( $\mu\text{M}$ ) in barley seedlings grown on agar or on floatable culture substrate.

		Agar	Floatable Substrate	
added concentration		Shoot	Shoot	Root
Hg 2+	0	2	1	54
	10	62	54	697
	20	171	144	851
	30	254	239	2060
	40	252	296	2430
Cd 2+	0	2	15	55
	10	15	4	479

	20	21	27	1270
	30	41	27	1450
	40	53	59	2880

The results show that uptake of  $\text{Hg}^{2+}$  and  $\text{Cd}^{2+}$  in barley shoots grown on floatable substrate or on agar medium is comparable. As expected there was a positive correlation between the metal-ion concentration in the medium and the concentration in the shoot tissue. The concentration of  $\text{Hg}^{2+}$  in the roots of seedlings grown on floatable substrate was increased by a factor of 10 compared to shoot tissue, and by a factor of 60 for  $\text{Cd}^{2+}$ .

Importantly, metal ion concentrations could not be established for root material grown on agar, because the roots could not be separated from the substrate material. Hence, metal ions contained in the substrate would have been measured in addition to those contained in the root material.

#### Example 3: Growing Arabidopsis or Thlaspi seedlings

A floatable substrate layer was prepared as described. Yet, particle size of the granulate was selected to comprise 1-5 mm, and a layer height of 3 cm was chosen. After adding MS-basal medium, seeds were directly applied to the upper surface of the layer. The seeds were germinated and cultivated for 21 days under conditions as described. At 21 days, the plants were harvested and placed on a dry filter paper to absorb attaching culture solution. The plants had reached a fresh weight of 0,5-2 g.

#### Example 4: Growing calluses on floatable culture substrate.

Tobacco internodes were surface sterilized as described. Floatable culture substrate and a MS-basal medium were filled into a glass culturing container such that a layer of approximately 3 cm height with a solution phase of equal height were formed. The glass container was lidded and autoclaved. After cooling, an internodal section of approximately 1 cm length was positioned on top of the substrate layer under sterile conditions. An appropriate amount of sterile IES and NAA stock solutions known to be useful for dedifferentiating tobacco tissue was added. The lid was sealed and the container placed in a controlled environment cabinet (relative humidity 70 +/- 5 %, 24 °C) and cultivated in the dark. After two weeks, callus tissue was separated from the internodal section under sterile conditions, was mechanically dissociated into smaller pieces and was placed in a new culturing container prepared as described. A representative picture of callus obtained after an additional week of culturing is shown in figure 3a, callus obtained after additional culturing for four weeks is shown in figure 3b.

**Example 5:** Growing meristems derived from tobacco plants on floatable culture substrate.

25

Tobacco plant shoots were harvested and surface sterilized as described. Under sterile conditions, the top two to three internodes of each shoot were dissected such that 6-8 leaves were preserved. The plant material had a height of approximately 30 mm and weighed approximately 5-7 g. Floatable culture substrate and a MS-basal medium were filled into a glass culturing container such that a layer of approximately 3 cm height with a free solution phase of

approximately 1 cm height were formed. The glass container was lidded and autoclaved. After cooling, the tobacco material was gently pushed into the substrate layer, taking care that none of the leaves was fully submerged (Figure 4a).

5 An appropriate amount of sterile IBA stock solution known to be useful for promoting root growth on tobacco tissue was added. After 10 days of culturing, the tobacco meristems showed richly developed primary roots and began to grow secondary roots (Figure 2b). The plants were supported by the  
10 substrate in an upright position, with notable growth of leave tissue (compare Figure 4a and 4b).

#### Example 6: Growing tobacco plants

15 A floatable culture layer with a height of 8 cm was prepared in a rectangular PE culturing vessel with a volume of 50 l. After adding approximately 20 l of MS basal medium to form a floating layer, the tobacco meristems of Example 5 were  
20 replanted in said layer at approximately 2 plants per 10 cm<sup>2</sup>. The meristems carrying rich primary roots were pushed into the substrate layer to a depth of approximately 2-2.5 cm, taking care that none of the leaves was fully submerged. The plants were grown under conditions as described for two  
25 weeks. The plants were harvested by gently pulling them out of the substrate layer. Adhering substrate material was detached by gently shaking the plants. At this point, the plants had reached an overall size of approximately 8-12 cm and a weight of 14-20 g. The shoots had grown to  
30 approximately 4-6 cm and had a normal morphology. The plants had a richly developed primary and secondary root system. The plants were replanted in conventional soil to be transferred to a greenhouse for hardening.

**Example 7:** Determination of the 'no adverse effect level (NOAEL)' for xenobiotics according to OECD guideline 208.

5

The plant of interest is selected. Amongst crop species e.g. the following can be used: tomato, cucumber, lettuce, soybean, cabbage, carrot, perennial ryegrass, corn or onion. A floatable culture substrate layer of appropriate height for the species of interest, typically 3-8 cm, is added to a culture vessel with a surface area of 15 cm<sup>2</sup>. MS basal medium or Hoaglands medium is prepared as described and added to the culture vessel such that a free solution phase of 1/3 to 1 x the height of the substrate layer is present. If appropriate, the vessel is lidded, autoclaved and left to cool prior to use.

The plant material is added at a plant density as follows:

One or two corn, soybean, tomato, cucumber or sugar beet plants per 15 cm container, three rape or pea plants per 15 cm container, 5-10 onion, wheat or other small seeds per 15 cm container. The number of containers is chosen to accommodate the planned range of different concentrations and replicate pots for each concentration of xenobiotic. A minimum of 20 plants per concentration, divided into a minimum of four replicates is required.

Stock solutions of the xenobiotic of interest are prepared and sterile-filtered, or autoclaved if appropriate.

Appropriate amounts of the stock solution are added to each of the containers to prepare a concentration range of the test substance (e.g. 0.1, 1.0, 10, 100 and 1000 mg / l medium). The test series also includes reference pots

containing no test substance, as well as at least one concentration of a different test substance with a known effect on the plant of interest. Solvent controls may be required for xenobiotics not dissolved in water. In every other respect, the control containers will be treated identical to the test containers. The test conditions should approximate those conditions necessary for normal growth for the species and varieties tested. To allow for defined culturing conditions, a growing chamber, phytotron, greenhouse etc. can be used. For the listed species the following conditions are recommended: carbon dioxide concentrations:  $350 \pm 50$  ppm, relative humidity:  $70 \pm 5$  % during light periods and  $90 \pm 5$  % during dark periods, temperature  $25 \pm 3$  °C during the day,  $20 \pm 3$  °C during the night, photoperiod: 16 h light / 8 h darkness, assuming an average wavelength of 400 to 700 nm, light: luminance of  $350 \pm 50$  micromol / m<sup>2</sup> / s, measured at the top of the canopy.

The plants are cultured for an observation period of 14-21 days after 50% of the control plants (also possible solvent controls) have emerged, the plants are observed frequently (at least weekly) for visual phytotoxicity and mortality. At the end of the test, measurement of % emergence and biomass should be recorded as well as visual phytotoxicity (chlorosis, necrosis, wilting, leaf and stem deformation).

For evaluation the plants are harvested by pulling them out of the substrate layer, gently shaking off adhering substrate particles and briefly rinsing with distilled water. Both the shoot and the root system can be evaluated. Regarding the root system, primary, and secondary roots as well as root hairs can be evaluated. Biomass can be measured using final shoot and root weight, preferably dry weight by harvesting

and drying at 60 °C to a constant weight. The results are recorded and evaluated using standard statistical procedures to calculate an EC 50, NOAEL etc.

## References

- Arteca RN and Arteca JM (2000) A novel method for growing  
5 Arabidopsis thaliana plants hydroponically. *Physiol Plant*  
108: 188-193.
- Feng J, Zheng SJ, Li XF, Takeda K and Matsumoto H (1997) A  
rapid hydroponic screening for aluminium tolerance in barley.  
10 *Plant Soil* 191: 133-137.
- Gibeaut DM, Hulett J, Cramer GR and Seemann JR (1997) Maximal  
biomass of Arabidopsis thaliana using a simple, low-  
maintenance hydroponic method and favourable environmental  
15 conditions. *Plant Physiol* 115: 317-319.
- Hart JJ, Welch RM, Norvell WA, Sullivan LA, Kochian LV (1998)  
Characterization of Cadmium Binding, Uptake, and  
Translocation in Intact Seedlings of Bread and Durum Wheat  
20 Cultivars. *Plant Physiol.* 116: 1413-1420.
- Hart JJ, Norvell WA, Welch RM, Sullivan LA, Kochian LV (1998)  
Characterization of Zinc Uptake, Binding and Translocation in  
Intact Seedlings of Bread and Durum Wheat Cultivars. *Plant*  
25 *Physiol.* 118: 219-226.
- Heidenreich B (1999) Transkriptionsänderungen von Arabidopsis  
thaliana bei Ozon-, UV-B und HgCl<sub>2</sub>-Behandlung. PhD thesis,  
LMU München.



Huttner D and Bar-Zvi D (2003) An improved, simple, hydroponic method for growing *Arabidopsis thaliana*. Plant Mol Biol Reprtr 21: 59-63.

- 5 Leal RS (2001) The use of Confidor S in the float, a new tobacco seedlings production system in the south of Brazil. Pflanzenschutz-Nachrichten Bayer 54, 3, 337-352.

- 10 Michalke B, Lustig S and Schramel P (1997) Analysis for the stability of platinum-containing species in soil samples using electrophoresis interfaced on-line with inductively coupled plasma mass spectrometry. Electrophoresis 18: 196-201.

- 15 Thiel S, Döhring T, Köfferlein M, Kosak A, Martin P and Seidlitz HK (1996) A phytotron for plant stress research: How far can artificial lighting compare to natural sunlight? J Plant Physiol 148: 456-463.

- 20 Tocquin P, Corbesier L, Havelange A, Pieltain A, Kurtem E, Bernier G and Périlleux C (2003) A novel high efficiency, low maintenance, hydroponic system for synchronous growth and flowering of *Arabidopsis thaliana*. BMC Plant Biol 3: 2.

Claims

1. A method for culturing plant material comprising:
  - (d) forming a layer of floatable granular substrate in  
5 a culturing vessel,
  - (e) placing plant material on or in said layer, and
  - (f) culturing the plant material in the presence of a culture medium, wherein there is no additional structure supporting the plant material from  
10 underneath.
2. A method according to Claim 1 wherein the culture medium is added before the layer of the granular substrate is formed.
3. A method according to Claim 1, wherein the culture  
15 medium is added after the layer of the granular substrate is formed.
4. A method according to any preceding Claim, wherein said floatable granular substrate comprises particles having an average diameter of 1-25 mm.
- 20 5. A method according to Claim 4, wherein said particles have a polygonal or spheroidal shape.
6. ~~A method according to Claim 4 or Claim 5, wherein said~~  
particles have an irregular polygonal or spheroidal shape.
- 25 7. A method according to Claim 4 or Claim 5, wherein said particles have a regular polygonal or spheroidal shape.

8. A method according to any preceding Claim, wherein said granular substrate comprises particles having a smooth surface.
- 5 9. A method according to any preceding Claim, wherein said granular substrate is chemically inert.
10. A method according to any preceding Claim, wherein said granular substrate has a density of 50 - 99.8 % of the density of the culture medium.
- 10 11. A method according to any preceding Claim, wherein said granular substrate is a thermoplastic polymer.
12. A method according to Claim 11, wherein said thermoplastic polymer is one or more of HD-PE, LD-PE or PP.
- 15 13. A method according to any preceding Claim, wherein said granular substrate has a density of 0.5 - 1.1 g/cm<sup>3</sup>.
14. A method according to Claim 13, wherein said granular substrate has a density of 0.90 - 0.96 g/cm<sup>3</sup>.
- 20 15. A method according to any preceding Claim, wherein said granular substrate comprises particles composed of more than one component, wherein said components can individually be more or less dense than the average density of the particle.
16. A method according to Claim 15, wherein said particles comprise at least one hollow enclosure.
- 25 17. A method according to any preceding Claim, further comprising an initial step of sterilizing the granular

substrate by a chemical treatment, irradiation and/or heat.

18. A method according to Claim 17, wherein said granular substrate is autoclaved.

5 19. A method according to any preceding Claim, wherein said granular substrate forms a substrate layer.

20. A method according to Claim 19, wherein said substrate layer is 0.5-20 cm. thick.

10 21. A method according to Claim 19 or Claim 20, wherein said substrate layer floats on the culture medium.

22. A method according to Claim 21, further comprising the step of aerating the culture medium.

15 23. A method according to any of Claims 19-22, wherein said substrate layer comprises additional embedded support structures.

24. A culturing kit for culturing plant material comprising a floatable granular culture substrate and a culturing vessel.

20 25. A culturing kit according to Claim 24, further comprising culturing solution and/or plant material.

-- 26. A method of culturing plant material comprising the use of a culturing kit according to Claim 24 or Claim 25.

27. Use of a floatable granular substrate for culturing plant material.

28. The use according to Claim 27, wherein said floatable granular substrate comprises particles having an average diameter of 1-25 mm.
- 5 29. The use according to Claim 28, wherein said particles have a polygonal or spheroidal shape
30. The use according to any of Claims 27-29, wherein said floatable granular substrate comprises particles having a smooth surface.
- 10 31. The use according to any of Claims 27-30, wherein said granular substrate is chemically inert.
32. The use according to any of Claims 27-31, wherein said granular substrate has a density of 50 - 99.8 % of the density of the culture medium used.
- 15 33. The use according to any of Claims 27-32, wherein said granular substrate is a thermoplastic polymer.
34. The use according to Claim 33, wherein said thermoplastic polymer is HD-PE, LD-PE, PP or a mixture of any thereof.
- 20 35. The use according to any of Claims 27-34, wherein said granular substrate has a density of 0.5 - 1.1 g/cm<sup>3</sup>.
36. The use according to Claim 35, wherein said granular substrate has a density of 0.90 - 0.96 g/cm<sup>3</sup>.
- 25 37. The use according to any of Claims 27-36, wherein said granular substrate comprises particles composed of more than one component, wherein said components can individually be more or less dense than the average density of the particle.

38. The use according to Claim 37, wherein said particles comprise at least one hollow enclosure.

**Abstract**

The present invention provides a method of culturing plant material comprising a layer of floatable granular substrate in a culturing vessel, plant material and culture medium.

- 5 Also, the invention provides a culturing kit comprising various combinations of floatable granular substrate with plant material, culturing solution and a culturing vessel, adapted to the specific requirements of hobby, science or industrial uses.




Figure 1






Figure 2a




Figure 2b




Figure 3a.






Figure 3b




Figure 4a




Figure 4b

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**